



SHORT REPORT

Open Access

Epidemiology of plasmid-mediated quinolone resistance in *salmonella enterica* serovar typhimurium isolates from food-producing animals in Japan

Tetsuo Asai^{1*}, Chizuru Sato², Kaori Masani², Masaru Usui¹, Manao Ozawa¹, Tomoe Ogino¹, Hiroshi Aoki², Takuo Sawada², Hidemasa Izumiya³, Haruo Watanabe³

Abstract

A total of 225 isolates of *Salmonella enterica* serovar Typhimurium from food-producing animals collected between 2003 and 2007 were examined for the prevalence of plasmid-mediated quinolone resistance (PMQR) determinants, namely *qnrA*, *qnrB*, *qnrC*, *qnrD*, *qnrS*, *qepA* and *aac(6)Ib-cr*, in Japan. Two isolates (0.8%) of *S. Typhimurium* DT104 from different dairy cows on a single farm in 2006 and 2007 were found to have *qnrS1* on a plasmid of approximately 9.6-kbp. None of the *S. Typhimurium* isolates had *qnrA*, *qnrB*, *qnrC*, *qnrD*, *qepA* and *acc(6)-Ib-cr*. Currently in Japan, the prevalence of the PMQR genes among *S. Typhimurium* isolates from food animals may remain low or restricted. The PFGE profile of two *S. Typhimurium* DT104 isolates without *qnrS1* on the farm in 2005 had an identical PFGE profile to those of two *S. Typhimurium* DT104 isolates with *qnrS1*. The PFGE analysis suggested that the already existing *S. Typhimurium* DT104 on the farm fortuitously acquired the *qnrS1* plasmid.

Findings

Salmonella enterica serovar Typhimurium is prevalent in many animal species [1-3] including food-producing animals that are considered to be reservoirs for human infection. *S. Typhimurium* was the top 5 serovar found most frequently in cases of *Salmonella* foodborne illness in Japan between 2006 and 2010 <https://hasseidoko.mhlw.go.jp/Byogentai/Pdf/data48e.pdf>. Multidrug-resistant *S. Typhimurium* definitive phage type 104 (DT104) causes human salmonellosis in Japan [3]. *S. Typhimurium* DT104 was first isolated in the late 1980 s, and has spread widely among food-producing animals across Japan [3-5]. Although a decreased proportion of DT104-related isolates among the animals was found between 2002 and 2005, multidrug-resistant *S. Typhimurium* remains prevalent among food-producing animals in Japan [6].

In Japan, fluoroquinolone drugs were approved in veterinary fields in 1991 and are commonly used for treatment of bacterial diseases such as enteritis and pneumonia in food-producing animals [7]. In 2001, fluoroquinolone resistance was found in *S. Choleraesuis* from pigs [8] and *S. Typhimurium* from cattle [9]. In addition, a fluoroquinolone-resistant *S. Typhimurium* was identified in bovine isolates in 2005 [6]. The mechanism of fluoroquinolone resistance in these isolates is the mutation of quinolone resistance-determining regions (QRDRs) in DNA gyrase and topoisomerase IV [8,9]. In 2006, *qnrS1* was identified in two *S. Typhimurium* isolates (including one DT104 isolate) from dairy cows and beef cattle, and *S. Thompson* from poultry in Japan [10]. The report identified the potential risk of foodborne infections of *Salmonella* conferring the gene from food-producing animals to humans in Japan.

Quinolone resistance mechanisms mediated by plasmids are responsible for target protection such as the *qnr* genes, active efflux such as *qepA*, and enzymatic modifications such as *aac(6)Ib-cr* [11]. The plasmid-mediated quinolone resistance (PMQR) genes contribute

* Correspondence: asai-t@nval.maff.go.jp

¹National Veterinary Assay Laboratory, Ministry of Agriculture, Forestry and Fisheries, 1-15-1 Tokura, Kokubunji, Tokyo 185-8511, Japan
Full list of author information is available at the end of the article

to a reduction of quinolone susceptibility. In Japan, *qnrS* was first identified in human isolates of *Shigella flexneri* in 2003 [12]. *qepA*-harboring clinical isolates of *Escherichia coli* were found in 2002 in Japan [13]. *qnrB* in *Klebsiella oxytoca*, *Pseudomonas mirabilis*, and *P. fluorescens*, and *qnrS* in *E. coli* and *Enterobacter cloacae* were found in zoo animal isolates in 2006 [14]. In addition, the presences of *qnrS1* and *qnrS2* in *Salmonella* isolated from fecal samples of overseas travelers were reported in Japan [15]. These reports provided an infectious source of *Enterobacteriaceae* conferring plasmid-mediated quinolone resistance in Japan. We examined the prevalence of plasmid-mediated quinolone resistance in *S. Typhimurium* isolated from food-producing animals.

A total of 225 isolates of *S. Typhimurium* from food-producing animals collected between 2003 and 2007 were derived from 156 cattle, 62 pigs and 7 poultry: includes 42 isolates of DT104, 8 of DT104B, and 2 of U302 (Table 1). Bacteriophage typing was performed according to the methods of the Health Protection Agency, London, United Kingdom [16]. Of the isolates, 132 *S. Typhimurium* isolates collected between 2003 and 2005 [6] were subjected to detection of the PMQR genes. The remaining 93 isolates between 2006 and 2007 were investigated for the presence of the PMQR genes and antimicrobial susceptibility. The presence of *qnrA*, *qnrB* and *qnrS* genes was determined by PCR [17]. The *qnrC* and *qnrD* genes were detected using the primers as previously described [18,19], respectively. The *qepA* and *acc(6')-Ib-cr* genes were examined as previously described [20,21]. Nucleotide sequences of both strands were determined directly on PCR products. The DNA alignments and deduced amino acid sequences were examined using the BLAST program (National Center for Biotechnology Information, USA). Minimum inhibitory concentrations (MICs) of antimicrobial agents were determined using the agar dilution methods according to the Clinical and Laboratory Standards Institute (CLSI) guidelines [22]. The following 11 antimicrobials were tested: ampicillin (ABPC), cefazolin, colistin, chloramphenicol (CP), dihydrostreptomycin

(DSM), gentamicin, kanamycin, oxytetracycline (OTC), nalidixic acid, enrofloxacin (ERFX), and trimethoprim. The MICs of each antimicrobial agent were interpreted using the recommendations of the CLSI [23]. The breakpoints not seen in the CLSI were defined in a previous study [1]. *Staphylococcus aureus* ATCC 29213, *Enterococcus faecalis* ATCC29212, *E. coli* ATCC 25922 and *P. aeruginosa* ATCC 27853 were used as quality control strains.

Of 225 *S. Typhimurium* isolates, two isolates of DT104, 18-PLS-16 and 19-PLS-45, from different dairy cows on a single farm in 2006 and 2007 showed *qnrS* positive results. The sequencing of amplicons showed complete identity to *qnrS1* previously identified on pAH0376 from a *S. flexneri* strain. None of the *S. Typhimurium* isolates had *qnrA*, *qnrB*, *qnrC*, *qnrD*, *qepA* and *acc(6')-Ib-cr*. The two isolates exhibited ERFX resistance (ERFX MIC, 2 mg/L) with resistances to ABPC, DSM, OTC and CP (Table 2).

The QRDR of *gyrA*, *parC* and *parE* was examined in ERFX-resistant isolates by PCR amplification and sequencing using primers as described elsewhere [24]. In addition, susceptibility of ERFX-resistant isolates to fluoroquinolones was examined using the micro broth dilution methods according to CLSI guidelines [22]. For evaluation of active efflux of the ERFX-resistant bacteria, the MIC of ERFX was determined by the agar dilution method in the presence of carbonyl cyanide *m*-chlorophenylhydrazone (CCCP) (100 μM). They had no mutations in the QRDR of GyrA, ParC and ParE. The MIC of ERFX was not changed in the presence of CCCP (100 μM). The two isolates with *qnrS1* exhibited almost the same MIC observed for each fluoroquinolone, which is relative low compared with the MIC for isolate (17-PLS-75) with mutations in the QRDR of GyrA and ParC.

Plasmid DNA was isolated from the *qnrS1*-positive isolates by the alkaline lysis method [25]. Extracted plasmids were transferred to Hybond-N+ membrane (Amersham Biosciences, Buckinghamshire, UK) using capillary blotting apparatus. The *qnrS1* PCR product was labeled with DIG-11-dUTP by PCR using a DIG High Prime DNA Labeling Kit (Roche Diagnostics Ltd,

Table 1 Salmonella Typhimurium isolates used in this study

Isolation year	Cattle				Pig			Poultry		
	Typhimurium	Phagetype			Typhimurium	Phagetype		Typhimurium	Phagetype	
		104	104B	U302		104	104B		104	
2003	24	8	2	0	8	0	0	0	0	
2004	25	3	0	2	8	1	0	0	0	
2005	42	12	0	0	21	1	0	4	1	
2006	23	4	0	0	11	2	0	2	1	
2007	42	4	4	0	14	5	2	1	0	
Total	156	31	6	2	62	9	2	7	2	

Table 2 Susceptibility for several fluoroquinolones

Antimicrobials	18-PLS-16	19-PLS-45	17-PLS-75
Year isolated	2006	2007	2005
Sources	Cattle	Cattle	Cattle
qnr	qnrS1	qnrS1	-
mutation in gyrA	WT	WT	S83F&D87N
mutation in parC	WT	WT	S80R
phagetype	104	104	12
Nalidixic acid	32	32	256
Oxolinic acid	4	4	>64
Flumequine	16	8	>64
Benofloxacin	4	4	16
Ciprofloxacin	1	1	8
Danofloxacin	2	2	16
Difloxacin	8	4	>32
Enrofloxacin	2	2	16
Levofloxacin	1	1	8
Norfloxacin	2	2	16

East Sussex, UK). After hybridization with the *qnrS1* probe, hybridized DNA was detected using a DIG Nucleic Acid Detection Kit (Roche Diagnostics Ltd). Using a plasmid profiling test, an approximately 93-kbp plasmid (virulence plasmid) was found in all four isolates, whereas there was also an approximately 9.6-kbp plasmid found in the *qnr*-conferring isolates. Hybridization tests revealed that *qnrS1* was located on the 9.6-Kbp plasmid (Figure 1).




Figure 1 Plasmid profile of *S.* Typhimurium isolates on a farm.
 Lane 1: 17-PLS-27, Lane 2: 17-PLS-28, Lane 3: 18-PLS-16, Lane 4: 19-PLS-45, Lane 5: Super coiled DNA ladder.

The appearance of *S.* Typhimurium DT104 conferring *qnrS1* on the farm is caused either by the introduction of *S.* Typhimurium DT104 conferring *qnrS1* or the transfer of the *qnrS1* plasmid to *S.* Typhimurium DT104 already existing on the farm. According to the CDC PulseNet protocol [26], genetic relatedness of isolates were analyzed by PFGE with XbaI and BlnI restriction enzymes. The isolates tested included two *qnrS1*-negative isolates of *S.* Typhimurium DT104 isolated in 2005 on a farm in which *qnrS1*-conferring isolates were found. In the present study, it was difficult to precisely distinguish between the two *S.* Typhimurium DT104 isolates without *qnrS1* and the two *S.* Typhimurium DT104 isolates with *qnrS1* by PFGE analysis (Figure 2). Our previous study showed that there is a variation in the BlnI-digested PFGE profiles of *S.* Typhimurium DT104 isolated from food-producing animals in Japan [5]. These results suggested that the *S.* Typhimurium DT104 already present on the farm fortuitously acquired the *qnrS1* plasmid. Previous studies showed that *qnrS1* in Typhimurium isolated in the UK was present on plasmids of 10,066 bp, which were transferable by the conjugation test and carry an IncN replicon [27,28]. Further study need to clarify the source of plasmid bearing *qnrS1*.




Figure 2 PFGE profile of *S.* Typhimurium isolates on a farm.
 Electrophoresis was performed using a CHEF-DR III System (Bio-Rad Laboratories, Hercules, CA, USA) with running conditions of 1 phase from 2.2 to 63.8 s at 180 V for 19 h. Lane1, 6, 11: *Salmonella* Braenderup H9812 digested with XbaI. Lane 2 and 7: 17-PLS-27, Lane 3 and 8: 17-PLS-28, Lane 4 and 9: 18-PLS-16, Lane 5 and 10: 19-PLS-45. Lane 2 to 5: XbaI digestion, Lane 7 to 10: BlnI digestion.

This study demonstrated that the two isolates of *S. Typhimurium* collected from different cattle on a farm in 2006 and 2007 harbored *qnrS1* on a 9.6-Kbp plasmid. At present in Japan, dissemination of *qnrS1* among *S. Typhimurium* isolates from food animals may remain restricted. The spread of plasmids carrying *qnr* among *Salmonella* isolates of animal origin could have serious consequences for fluoroquinolone treatment of non-typhoid *Salmonella* infection in humans and animals. Previously, *qnrS1* and *qnrS2* were found in serovars Typhimurium, Corvallis, Montevideo, Agona, Braenderup and Alacua of *Salmonella* isolates from fecal samples of overseas travelers who had visited Thailand, Malaysia, Vietnam, Indonesia and Singapore, between 2001 and 2007 [15]. PMQR is identified in human isolates of *Enterobacteriaceae* but is likely to be rare in isolates from food-producing animals [29]. However, in China, plasmid-mediated quinolone resistance is frequently found in the isolates from food-producing animals [20]. Thus it would be difficult to prevent the invasion of resistance genes from foreign countries to Japan. The monitoring of fluoroquinolone use and quinolone resistance in bacteria of food-producing animal origin is essential to assess the level of risk of resistance in food-borne bacteria in the animals.

Abbreviations

ABPC: ampicillin; CCCP: carbonyl cyanide *m*-chlorophenylhydrazone; CLSI: Clinical and Laboratory Standards Institute; CP: chloramphenicol; DSM: dihydrostreptomycin; DT104: definitive phage type 104; MICs: Minimum inhibitory concentrations OTC: oxytetracycline; PMQR: plasmid-mediated quinolone resistance; QRDRs: quinolone resistance-determining regions; ERFX: enrofloxacin.

Acknowledgements

We thank the staff of the Livestock Hygiene Service Centers across Japan for providing *S. enterica* serovar Typhimurium isolates. This work was supported in part by a grant-in aid from the Japanese Ministry of Health, Labour and Welfare (H21-Shokuhin-Ippan-013).

Author details

¹National Veterinary Assay Laboratory, Ministry of Agriculture, Forestry and Fisheries, 1-15-1 Tokura, Kokubunji, Tokyo 185-8511, Japan. ²Nippon Veterinary and Life Science University, 1-7-1 Kyonancho, Musashino, Tokyo 180-8602, Japan. ³National Institute of Infectious Diseases, 1-23-1 Toyama, Shinjuku-ku, Tokyo 162-8640, Japan.

Authors' contributions

TA conceived the study, the study design, participated in the determination of quinolone resistance and determinants, interpreted the data and drafted the manuscript. CS carried out large parts of the experimental work. KM helped to carried out prevalence of resistance genes. MU helped to carried out prevalence of resistance genes. MO carried out the antimicrobial susceptibility testing. TO carried out the antimicrobial susceptibility testing. HA helped to carry out determination of quinolone resistance and draft the manuscript. TS helped to draft the manuscript. HI carried out phage typing and helped to draft the manuscript. WH helped to draft the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 17 November 2010 Accepted: 7 December 2010
Published: 7 December 2010

References

1. Asai T, Esaki H, Kojima A, Ishihara K, Tamura Y, Takahashi T: Antimicrobial resistance in *Salmonella* isolates from apparently healthy food-producing animal from 2000 to 2003: the first stage of Japanese veterinary antimicrobial resistance monitoring (JVARM). *J Vet Med Sci* 2006, **68**:881-884.
2. Asai T, Otagiri Y, Osumi T, Namimatsu T, Hirai H, Sato S: Isolation of *Salmonella* from diarrheic feces of pigs. *J Vet Med Sci* 2002, **64**:159-160.
3. Izumiya H, Tamura K, Terajima J, Watanabe H: *Salmonella enterica* serovar Typhimurium phage type DT104 and other multi-drug resistant strains in Japan. *Jpn J Infect Dis* 1999, **52**:133.
4. Sameshima T, Akiba M, Izumiya H, Terajima J, Tamura K, Watanabe H, Nakazawa M: *Salmonella typhimurium* DT104 from livestock in Japan. *Jpn J Infect Dis* 2000, **53**:15-16.
5. Esaki H, Morioka A, Kojima A, Ishihara K, Asai T, Tamura Y, Izumiya H, Terajima J, Watanabe H, Takahashi T: Epidemiological characterization of *Salmonella Typhimurium* DT104 prevalent among food-producing animals in the Japanese veterinary antimicrobial resistance monitoring program (1999-2001). *Microbiol Immunol* 2004, **48**:553-556.
6. Kawagoe K, Mine H, Asai T, Kojima A, Ishihara K, Harada K, Ozawa M, Izumiya H, Terajima J, Watanabe H, et al: Changes of multi-drug resistance pattern in *Salmonella enterica* subspecies *enterica* serovar Typhimurium isolates from food-producing animals in Japan. *J Vet Med Sci* 2007, **69**:1211-1213.
7. Asai T, Harada K, Ishihara K, Kojima A, Sameshima T, Tamura Y, Takahashi T: Association of antimicrobial resistance in *Campylobacter* isolated from food-producing animals with antimicrobial use on farms. *Jpn J Infect Dis* 2007, **60**:290-294.
8. Esaki H, Chiu CH, Kojima A, Ishihara K, Asai T, Tamura Y, Takahashi T: Comparison of fluoroquinolone resistance genes of *Salmonella enterica* serovar Choleraesuis isolates in Japan and Taiwan. *Jpn J Infect Dis* 2004, **57**:287-288.
9. Izumiya H, Terajima J, Matsushita S, Tamura K, Watanabe H: Characterization of multidrug-resistant *Salmonella enterica* serovar Typhimurium isolated in Japan. *J Clin Microbiol* 2001, **39**:2700-2703.
10. Ahmed AM, Ishida Y, Shimamoto T: Molecular characterization of antimicrobial resistance in *Salmonella* isolated from animals in Japan. *J Appl Microbiol* 2009, **106**:402-409.
11. Jacoby GA: Mechanisms of resistance to quinolones. *Clin Infect Dis* 2005, **41**(Suppl 2):S120-126.
12. Hata M, Suzuki M, Matsumoto M, Takahashi M, Sato K, Ibe S, Sakae K: Cloning of a novel gene for quinolone resistance from a transferable plasmid in *Shigella flexneri* 2b. *Antimicrob Agents Chemother* 2005, **49**:801-803.
13. Yamane K, Wachino J, Suzuki S, Kimura K, Shibata N, Kato H, Shibayama K, Konda T, Arakawa Y: New plasmid-mediated fluoroquinolone efflux pump, QepA, found in an *Escherichia coli* clinical isolate. *Antimicrob Agents Chemother* 2007, **51**:3354-3360.
14. Ahmed AM, Motoi Y, Sato M, Maruyama A, Watanabe H, Fukumoto Y, Shimamoto T: Zoo animals as reservoirs of gram-negative bacteria harboring integrons and antimicrobial resistance genes. *Appl Environ Microbiol* 2007, **73**:6686-6690.
15. Taguchi M, Kawahara R, Seto K, Inoue K, Hayashi A, Yamagata N, Kamakura K, Kashiwagi E: Plasmid-mediated quinolone resistance in *Salmonella* isolated from patients with overseas travelers' diarrhea in Japan. *Jpn J Infect Dis* 2009, **62**:312-314.
16. Anderson ES, Ward LR, Saxe MJ, de Sa JD: Bacteriophage-typing designations of *Salmonella typhimurium*. *J Hyg (Lond)* 1977, **78**:297-300.
17. Cattoir V, Poirel L, Rotimi V, Soussy CJ, Nordmann P: Multiplex PCR for detection of plasmid-mediated quinolone resistance *qnr* genes in ESBL-producing enterobacterial isolates. *J Antimicrob Chemother* 2007, **60**:394-397.
18. Wang M, Guo Q, Xu X, Wang X, Ye X, Wu S, Hooper DC, Wang M: New plasmid-mediated quinolone resistance gene, *qnrC*, found in a clinical isolate of *Proteus mirabilis*. *Antimicrob Agents Chemother* 2009, **53**:1892-1897.

19. Cavaco LM, Hasman H, Xia S, Aarestrup FM: **qnrD, a novel gene conferring transferable quinolone resistance in *Salmonella enterica* serovar Kentucky and Bovismorbificans strains of human origin.** *Antimicrob Agents Chemother* 2009, **53**:603-608.
20. Ma J, Zeng Z, Chen Z, Xu X, Wang X, Deng Y, Lu D, Huang L, Zhang Y, Liu J, et al: **High prevalence of plasmid-mediated quinolone resistance determinants qnr, aac(6)-Ib-cr, and qepA among ceftiofur-resistant *Enterobacteriaceae* isolates from companion and food-producing animals.** *Antimicrob Agents Chemother* 2009, **53**:519-524.
21. Park CH, Robicsek A, Jacoby GA, Sahn D, Hooper DC: **Prevalence in the United States of aac(6)-Ib-cr encoding a ciprofloxacin-modifying enzyme.** *Antimicrob Agents Chemother* 2006, **50**:3953-3955.
22. CLSI: **Performance standards for antimicrobial disk and dilution susceptibility tests for bacteria isolated from animals.** *Approved standard M31-A3*. 3 edition. Wayne, PA: Clinical and Laboratory Standards Institute; 2008.
23. CLSI: **Performance standards for antimicrobial susceptibility testing; eighteenth informational supplement.** Wayne, PA: Clinical and Laboratory Standards Institute; 2008.
24. Giraud E, Brisabois A, Martel JL, Chaslus-Dancla E: **Comparative studies of mutations in animal isolates and experimental in vitro- and in vivo-selected mutants of *Salmonella* spp. suggest a counterselection of highly fluoroquinolone-resistant strains in the field.** *Antimicrob Agents Chemother* 1999, **43**:2131-2137.
25. Birnboim HC, Doly J: **A rapid alkaline extraction procedure for screening recombinant plasmid DNA.** *Nucleic Acids Res* 1979, **7**:1513-1523.
26. Hunter SB, Vauterin P, Lambert-Fair MA, Van Duyn MS, Kubota K, Graves L, Wrigley D, Barrett T, Ribot E: **Establishment of a universal size standard strain for use with the PulseNet standardized pulsed-field gel electrophoresis protocols: converting the national databases to the new size standard.** *J Clin Microbiol* 2005, **43**:1045-1050.
27. Kehrenberg C, Hopkins KL, Threlfall EJ, Schwarz S: **Complete nucleotide sequence of a small qnrS1-carrying plasmid from *Salmonella enterica* subsp. *enterica* Typhimurium DT193.** *J Antimicrob Chemother* 2007, **60**:903-905.
28. Hopkins KL, Wootton L, Day MR, Threlfall EJ: **Plasmid-mediated quinolone resistance determinant qnrS1 found in *Salmonella enterica* strains isolated in the UK.** *J Antimicrob Chemother* 2007, **59**:1071-1075.
29. Robicsek A, Jacoby GA, Hooper DC: **The worldwide emergence of plasmid-mediated quinolone resistance.** *Lancet Infect Dis* 2006, **6**:629-640.

doi:10.1186/1757-4749-2-17

Cite this article as: Asai et al.: Epidemiology of plasmid-mediated quinolone resistance in *salmonella enterica* serovar typhimurium isolates from food-producing animals in Japan. *Gut Pathogens* 2010 **2**:17.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

